

First Name: \_\_\_\_\_ Last Name: \_\_\_\_\_

Student-No: \_\_\_\_\_ Section: \_\_\_\_\_

**You must show your work**

1. 3 marks A wholesale flour producer wants to sell flour in cubic containers. The cost of making a cubic container with edge  $x$  meters is  $3x^3 + 87x^{\frac{3}{2}}$  dollars. A market research shows that people are willing to be  $y$  cubic meters of flour for  $91\sqrt{y}$  dollars. Find the size of the edge of the box which will produce the maximal profit per box.

Answer:  $x = \left(\frac{5}{6}\right)^{\frac{2}{3}}$

**Solution:** The volume of the box is  $V(x) = x^3$  and the revenue per box is

$$R(x) = 91\sqrt{V(x)} = 91x^{\frac{3}{2}}.$$

The profit function is

$$P(x) = R(x) - C(x) = 4x^{\frac{3}{2}} - 3x^3.$$

We are interested in values of  $x$  in the open interval  $(0, \infty)$ . We look for critical points

$$P'(x) = \frac{12}{2}\sqrt{x} - 9x^2 = 6\sqrt{x} - 9x^2$$

Solving  $P'(x) = 0$  for  $x$  we get  $x^{\frac{3}{2}} = \frac{2}{3}$  so the only critical point of  $P(x)$  is at  $\left(\frac{2}{3}\right)^{\frac{2}{3}}$ .

The second derivative of  $P$  is

$$P''(x) = \frac{3}{\sqrt{x}} - 27x.$$

Which is positive for  $0 < x < \left(\frac{1}{9}\right)^{\frac{2}{3}}$  and negative for  $x > \left(\frac{1}{9}\right)^{\frac{2}{3}}$ . In particular, it follows that  $P$  is increasing for  $0 < x < \left(\frac{2}{3}\right)^{\frac{2}{3}}$  and decreasing for  $x > \left(\frac{2}{3}\right)^{\frac{2}{3}}$  so  $x = \left(\frac{2}{3}\right)^{\frac{2}{3}}$  is the absolute maximum of  $P(x)$  at  $(0, \infty)$ . The profit is going to be \$0.75.

2. 3 marks Approximate  $\sqrt[3]{15}$  as rational number, using a 2-nd Taylor Polynomial of  $f(x) = \sqrt[3]{x}$ .

Answer:  $2 + \frac{7}{12} - \frac{7^2}{288}$  or  $3 - \frac{12}{27} - \frac{12^2}{2187}$

**Solution:**

- The closest cubes to 15 are 8 and 27. I solve for both choices but you needed choose only one. Write

$$A(x) = a_0 + a_1(x - 8) + a_2(x - 8)^2$$

$$B(x) = b_0 + b_1(x - 27) + b_2(x - 27)^2$$

- 0-term:

$$a_0 = f(8) = 2, \quad b_0 = f(27) = 3.$$

In particular  $2 < \sqrt[3]{15} < 3$  (since  $f$  is increasing).

- 1-term:  $f'(x) = \frac{1}{3x^{\frac{2}{3}}}$

$$a_1 = f'(8) = \frac{1}{3 \cdot 2} = \frac{1}{12}, \quad b_1 = f'(27) = \frac{1}{3 \cdot 9} = \frac{1}{27}$$

- 2-term:  $f''(x) = \frac{-2}{9x^{\frac{5}{3}}}$

$$a_2 = \frac{f''(8)}{2} = \frac{-2}{2 \cdot 9 \cdot 2^5} = \frac{-1}{288}, \quad b_2 = \frac{f''(27)}{2!} = \frac{-2}{2 \cdot 9 \cdot 243} = \frac{-1}{2187}$$

- The 2-nd taylor polynomial centered at  $x = 8$  is

$$A(x) = 2 + \frac{1}{12}(x - 8) - \frac{1}{288}(x - 8)^2$$

$$B(x) = 3 + \frac{1}{27}(x - 27) - \frac{1}{2187}(x - 27)^2$$

- The approximations of  $\sqrt[3]{15}$  are given by

$$A(15) = 2 + \frac{7}{12} - \frac{7^2}{288}, \quad B(15) = 3 - \frac{12}{27} - \frac{12^2}{2187}.$$

3. 4 marks Sketch the graph of the function  $f(x) = \frac{(x-16)^2}{x^2-16}$ . You may use the fact that  $f''(x) = 0$  only at  $x_0 = \frac{17+5\sqrt[3]{45}+3\sqrt[3]{75}}{2} \approx 23.7$ .

### Solution:

- The domain of  $f(x)$  is  $x \neq \pm 4$ , i.e.  $(-\infty, -4)$ ,  $(-4, 4)$  and  $(4, \infty)$ .
- Vertical asymptotes:

$$\lim_{x \rightarrow 4^+} f(x) = \infty$$

$$\lim_{x \rightarrow 4^-} f(x) = -\infty$$

$$\lim_{x \rightarrow -4^+} f(x) = -\infty$$

$$\lim_{x \rightarrow -4^-} f(x) = \infty$$

Horizontal asymptotes:

$$\lim_{x \rightarrow \infty} f(x) = 1$$

$$\lim_{x \rightarrow -\infty} f(x) = \lim_{x \rightarrow \infty} f(-x) = 1$$

- The first derivative of  $f(x)$  is

$$f'(x) = \frac{32(x-16)(x-1)}{(x^2-16)^2}$$

The derivative is defined for any  $x$  in the domain (i.e.  $x \neq \pm 4$ ) and so the only critical points are the points  $c$  where  $f'(c) = 0$ , i.e.  $c = 1$  and  $c = 16$ . Also, since the denominator is always positive, the sign of the first derivative is the same as the sign of its numerator. So,  $f'(x)$  is positive (and  $f(x)$  is increasing) for  $x < -4$  and  $x > 4$  and it is negative (and  $f(x)$  is decreasing) for  $-4 < x < 4$ . It follows that  $c = 1$  is a local maximum and  $c = 16$  is a local minimum.

- $f(1) = -15$ ,  $f(16) = 0$
- The second derivative of  $f(x)$  is

$$f''(x) = 32 \frac{-2x^3 + 51x^2 - 96x + 272}{(x^2 - 9)^3}$$

The denominator is positive for  $x > 4$  and  $x < -4$  and negative for  $-4 < x < 4$ . The numerator is negative for  $x > x_0$  and positive for  $x < x_0$ . It follows that  $f''(x)$  is positive (and  $f(x)$  is concave up) for  $x < -4$  and  $3 < x < x_0$  and negative (and  $f(x)$  is concave down) for  $-4 < x < 4$  and  $x > x_0$ . So  $x_0$  is an inflection point.

