

Linear Approximation

Nov 29
Lec 24


Local Linearization

Approximate a function (usually complicated) with a line.

→ What is this line? The tangent line

Recall

→ Tangent line: The function $f(x)$ and a point $(a, f(a))$ on f are given



$(a, f(a))$ is the touch point, both on the graph and on the line. To write the equation of the tangent line we need:

- slope $\rightarrow m_{\text{tan}} = f'(a)$
- & point $\rightarrow (a, f(a))$

$$\Rightarrow y - f(a) = f'(a)(x - a)$$


or

$$y = \underbrace{f'(a)(x - a) + f(a)}_{\text{call this } L(x)} \rightarrow \text{tangent line equation}$$

then $L(x) = f'(a)(x - a) + f(a)$ is the tangent line equation.

Linear Approximation : If f is differentiable at "a" and x is a point close to "a", then the function $f(x)$ can be approximated with its tangent line $L(x)$.

Or $f(x)$ is close to $L(x) \rightsquigarrow f(x) \approx L(x)$



This is an example of over-estimate because the value of $L(x)$ is greater than $f(x)$.
 \downarrow
 $L(x)$ sits above $f(x)$.

$$f(x) \approx L(x)$$

$$L(x) = f'(a)(x-a) + f(a)$$

* Instead of evaluating f at x , we evaluate L at x and estimate f with L .

* In linearization questions, we should first find:

"a" : The touch point \rightsquigarrow The "good" point usually with no decimal points.

"x" : The point close to "a" \rightsquigarrow usually has decimals.

f : The function that we should use to find the linearization. Sometimes it's explicitly given in the question, but sometimes we should find it from the given estimate.

Example . a) Find the linear approximation of $f(x) = \sqrt{x}$ at $x=9$

tangent line at $x=9$

Find the touch point $x=9 \rightsquigarrow f(9) = \sqrt{9} = 3 \rightsquigarrow (9, 3)$

slope $\rightsquigarrow f'(9)$

$$f(x) = x^{\frac{1}{2}} \Rightarrow f'(x) = \frac{1}{2} x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(9) = \frac{1}{2\sqrt{9}} = \frac{1}{6}$$

$$\Rightarrow y - 3 = \frac{1}{6}(x - 9)$$

$$\Rightarrow L(x) = \frac{1}{6}(x - 9) + 3 \rightarrow \text{linear approximation of } f(x) = \sqrt{x}$$

b) Sketch the graph of f and its local linearization.


$$\rightsquigarrow f(x) = \sqrt{x}$$

$$f(0) = \sqrt{0} = 0$$

$$f(1) = \sqrt{1} = 1$$

$$f(4) = \sqrt{4} = 2$$

$$f(9) = \sqrt{9} = 3$$



$$\rightsquigarrow L(x) = \frac{1}{6}(x - 9) + 3$$

$$f(9) = 3$$

$$f(6) = \frac{1}{6}(6 - 9) + 3$$

$$= -\frac{0.5}{2} + 3$$

$$= 2.5$$

$(9, 3)$ is both
on the line and
on $f(x) = \sqrt{x}$

The tangent line always on the touch point, we don't change the tangent line, we use the same line to evaluate and estimate for x 's close to the touch point " a ".

c) find $\sqrt{9.1}$ and $\sqrt{8.88}$.

First find : "a", "x" and $f(x)$.

$f(x)$ is given : $f(x) = \sqrt{x}$

$\sqrt{9.1} = f(9.1) \rightsquigarrow \underset{\downarrow}{9.1}$ is close to $\underset{\downarrow}{9}$
"x" : the point close to "a" usually with decimal points. "a" : the "nice" point with no decimal points.

$$\Rightarrow f(x) \approx L(x)$$

$$\Rightarrow \sqrt{9.1} = f(9.1) \approx L(9.1)$$

from part (a) : $L(x) = \frac{1}{6}(x-9) + 3$

$$\Rightarrow L(9.1) = \frac{1}{6}(9.1-9) + 3 = \frac{0.1}{6} + 3 = 3.0166$$

$$\Rightarrow \boxed{\sqrt{9.1} \approx 3.0166}$$

Similarly :

$$\sqrt{8.88} = f(8.88) \approx L(8.88)$$

$$= \frac{1}{6}(8.88-9) + 3$$

$$= \frac{-0.12}{6} + 3$$

$$= -0.02 + 3 = 2.98$$

$$\Rightarrow \boxed{\sqrt{8.88} \approx 2.98}$$

Example Estimate $\ln(1.2)$ and $\ln(e^{-0.05})$

Find f , "a" and "x".

f is NOT given in the question, but since we are looking for "ln" of some number so: $f(x) = \ln x$

"a" is the "nice" touch point: $a = 1$

"x" is the point close to "a": $x = 1.2$

⇒ We use linear approximation to estimate the values, so first we need to find the equation to the line.

$$f(x) = \ln x$$

$$a = 1 \Rightarrow f(1) = \ln 1 = 0 \Rightarrow (1, 0) \text{ touch point}$$

$$f'(x) = \frac{1}{x} \Rightarrow f'(1) = \frac{1}{1} \Rightarrow m_{\text{tan}} = 1$$

$$\Rightarrow y - 0 = 1(x - 1)$$

$$\Rightarrow \boxed{L_1(x) = x - 1}$$

1.2 is close to 1, so we can use $L_1(x)$ to estimate:

$$\ln(1.2) = f(1.2) \approx L_1(1.2) = 1.2 - 1 = 0.2$$

$$\Rightarrow \boxed{\ln(1.2) \approx 0.2}$$

$\ln(e - 0.05)$: We can NOT use the previous $L(x)$, because the point "a" is different.

Again : $f(x) = \ln x$, But $a = e$ and $x = e - 0.05$

touch point
 $a = e$

$$\underbrace{f(e)}_{\ln e = 1} \Rightarrow (e, 1)$$

slope : $f'(e) = \frac{1}{e} \Rightarrow y - 1 = \frac{1}{e}(x - e)$

$$\Rightarrow L_2(x) = \frac{1}{e}(x - e) + 1$$

$e - 0.05$ is close to e , we use $L_2(x)$ to estimate :

$$\ln(e - 0.05) = f(e - 0.05) \approx L_2(e - 0.05)$$

$$= \frac{1}{e}(\cancel{e} - 0.05 - \cancel{e}) + 1$$

$$= \frac{-0.05}{e} + 1$$

Example. a) Find the linearization to the function

$$g(x) = x f(x^2) \quad \text{at } x=2$$

Given that $f(2) = -1$, $f'(2) = 6$, $f(4) = 3$, $f'(4) = -4$

b) estimate $g(1.99)$

a) linearization: tangent line at $x=2$

touch point: $g(2) = 2 f(2^2) = 2 f(4) = 2 \times 3 = 6 \rightarrow (2, 6)$

slope: $g'(2)$

$$g'(x) = \underbrace{1 \times f(x^2)}_{\text{product Rule}} + \underbrace{x \times f'(x^2)}_{\text{Chain rule}} \times \underbrace{2x}_{\text{inside}}$$

$$\begin{aligned} \Rightarrow g'(2) &= f(2^2) + 2 f'(2^2) \times 2 \times 2 \\ &= 3 + 2 \times -4 \times 4 = 3 - 32 = -29 \end{aligned}$$

$$\Rightarrow y - 6 = -29(x - 2) \Rightarrow L(x) = -29(x - 2) + 6$$

b) 1.99 is close to 2 \rightarrow Use $L(x)$ to estimate:

$$\begin{aligned} g(1.99) &\approx L(1.99) = -29(1.99 - 2) + 6 \\ &= -29 \times -0.01 + 6 \\ &= +0.29 + 6 \\ &= \underline{6.29} \end{aligned}$$

I'd rather NOT to distribute -29 because when evaluating $L(1.99)$ computations will be easier.

Practice . Use local linearization to estimate

(a) $(2.02)^8$

(b) $\sqrt[3]{0.98}$

(c) $e^{0.06}$

(a) $(2.02)^8$ what is f , a and x ?

something is raised to power 8 $\Rightarrow f(x) = x^8$

$a = 2$ the "nice" touch point

$x = 2.02$

$f(x) = x^8 \Rightarrow f(2) = 2^8 \Rightarrow (2, 2^8)$ touch point

$f'(x) = 8x^7 \Rightarrow f'(2) = 8 \cdot 2^7 = m_{\text{tan}}$

$\Rightarrow y - 2^8 = 8 \cdot 2^7 (x - 2)$

$\Rightarrow L(x) = 8 \cdot 2^7 (x - 2) + 2^8$

2.02 is close to 2, we use $L(x)$ to estimate:

$(2.02)^8 = f(2.02) \approx L(2.02)$

$= 8 \cdot 2^7 (2.02 - 2) + 2^8$

$= 2^3 \cdot 2^7 \cdot 0.02 + 2^8$

$= 20.48 + 256 = \underline{276.48}$

(b) $\sqrt[3]{0.98} \rightsquigarrow$ taking the cubic root $\rightsquigarrow f(x) = \sqrt[3]{x}$
 $a = 1$, $x = 0.98$

$f(x) = x^{\frac{1}{3}} \Rightarrow f(1) = \sqrt[3]{1} = 1 \rightarrow (1, 1)$ touch point

$f'(x) = \frac{1}{3} x^{\frac{1}{3}-1} = \frac{1}{3} x^{-\frac{2}{3}} = \frac{1}{3 x^{\frac{2}{3}}} = \frac{1}{3 \sqrt[3]{x^2}}$

$\Rightarrow f'(1) = \frac{1}{3} \Rightarrow y - 1 = \frac{1}{3} (x - 1)$

$\Rightarrow \underline{L(x) = \frac{1}{3}(x-1) + 1}$

0.98 is close to $1 \Rightarrow \sqrt[3]{0.98} = f(0.98) \approx L(0.98)$

$L(0.98) = \frac{1}{3} (0.98 - 1) + 1 = \underline{\underline{-\frac{0.02}{3} + 1 \approx \sqrt[3]{0.98}}}$

(c) $e^{0.06} \rightsquigarrow f(x) = e^x$, 0.06 is close to 0 , therefore
 $a = 0$ and $x = 0.06$

$f(0) = e^0 = 1 \Rightarrow (0, 1)$ touch point

$f'(x) = e^x \Rightarrow m_{\tan} = f'(0) = e^0 = 1 \rightsquigarrow y - 1 = 1(x - 0)$

$\Rightarrow \underline{L(x) = x + 1}$

$e^{0.06} = f(0.06) \approx L(0.06) = 0.06 + 1 = 1.06$

$\Rightarrow \underline{e^{0.06} \approx 1.06}$